

Deeply virtual electro-production of vector mesons and spin effects

S.V.GOLOSKOKOV

*Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research,
Dubna, Russia*

We study light vector meson photoproduction at small x on the basis of the generalized parton distribution (GPD). Our results on the cross section and spin density matrix elements (SDME) are in fair agreement with DESY experiments.

PACS: 12.38.Bx, 13.60.Hb

Key words: Diffraction, spin, parton distributions, vector meson production

1 Introduction

Vector meson leptonproduction at large energies [1] is one of the important processes which is sensitive to the GPDs. At small x -Bjorken (x) the process factorizes [2] into a hard meson leptonproduction off gluons and GPD (Fig.1).




Fig. 1. The handbag diagram for the meson electroproduction off proton.

The generally used standard collinear approximation (SCA) leads to some essential problems. The cross section for longitudinally polarized virtual photons which predominates for the large photon virtualities exceeds the data by a large factor if calculated in the SCA [3]. Only the amplitude with a longitudinally polarized virtual photon and vector meson (LL) amplitude can be calculated in the SCA. The transition amplitudes for transversally polarized photons $\gamma_{\perp}^* \rightarrow V_{\perp}$ (TT) and $\gamma_{\perp}^* \rightarrow V_L$ (TL) exhibit in the SCA the infrared singularities [4] which breakdown factorization.

Here we use the modified perturbative approach (MPA) [5] which includes the quark transverse degrees of freedom accompanied by Sudakov suppressions. In the MPA one can solve all mentioned problems. Transverse quark degrees of freedom and Sudakov factors suppress the contribution from the end-point region in the LL amplitude. The transverse quark momentum regularizes the end-point singularities

in TT and TL amplitudes. This leads to reasonable agreement of our results with H1 and ZEUS data [6, 7] for electroproduced ρ and ϕ mesons at small x [1, 8].

In this report, we discuss the spin effects in the ρ meson leptonproduction. Within the MPA we calculate the three LL , TT and TL transition amplitudes, and afterwards cross sections and SDME.

2 Leptonproduction of Vector Mesons in the GPD approach

The leading-twist wave function $\hat{\Psi}_V^0$ describes the longitudinally polarized vector mesons. To study the transversally polarized light mesons, we use the higher order k -dependent wave function $\hat{\Psi}_V^1$ proposed in [9]

$$\hat{\Psi}_V = \hat{\Psi}_V^0 + \hat{\Psi}_V^1. \quad (1)$$

$$\hat{\Psi}_V^0 = (\not{V} + m_V) \not{\epsilon}_V \phi_V(k, \tau). \quad (2)$$

$$\hat{\Psi}_V^1 = [\frac{2}{M_V} \not{V} \not{K} - \frac{2}{M_V} (\not{V} - m_V)(\epsilon_V \cdot K)] \phi'_V(k, \tau). \quad (3)$$

Here ϵ_V is a polarization of the vector meson with momentum V , τ is the fraction of the meson momentum carried by the quark, M_V is the scale in the higher order wave function which should be about the meson mass m_V . We use here $M_V = m_V/2$

The gluon contribution to the leptonproduction amplitudes for $t \sim 0$ and positive proton helicity reads as a convolution of the hard subprocess amplitude $\mathcal{H}^V$ and a large distance unpolarized H^g and polarized $\tilde{H}^g$ gluon GPDs (Fig.1):

$$\begin{aligned} \mathcal{M}_{\mu'+, \mu+} &= \frac{e}{2} \mathcal{C}_V \int_0^1 \frac{d\bar{x}}{(\bar{x} + \xi)(\bar{x} - \xi + i\varepsilon)} \\ &\times \left\{ \left[\mathcal{H}_{\mu'+, \mu+}^V + \mathcal{H}_{\mu'-, \mu-}^V \right] H^g(\bar{x}, \xi, t) \right. \\ &\left. + \left[\mathcal{H}_{\mu'+, \mu+}^V - \mathcal{H}_{\mu'-, \mu-}^V \right] \tilde{H}^g(\bar{x}, \xi, t) \right\}. \end{aligned} \quad (4)$$

Here μ (μ') denotes the helicity of the photon (meson), $\bar{x}$ is the momentum fraction of the transversally polarized gluons and the skewness ξ is related to Bjorken- x by $\xi \simeq x/2$. The flavor factors are $C_\rho = 1/\sqrt{2}$ and $C_\phi = -1/3$. The polarized $\tilde{H}^g$ GPD at small $\bar{x}$ is much smaller than the unpolarized GPD H^g and will be important in the A_{LL} asymmetry only.

The subprocess amplitude $\mathcal{H}^V$ is represented as the contraction of the hard part F , which is calculated perturbatively, and the non-perturbative meson wave function ϕ_V

$$\mathcal{H}_{\mu'+, \mu+}^V \pm \mathcal{H}_{\mu'-, \mu-}^V = \frac{2\pi\alpha_s(\mu_R)}{\sqrt{2N_c}} \int_0^1 d\tau \int \frac{d^2\mathbf{k}_\perp}{16\pi^3} \phi_V(\tau, k_\perp^2) F_{\mu'\mu}^\pm. \quad (5)$$

The wave function is chosen in Gaussian form

$$\phi_V(\mathbf{k}_\perp, \tau) = 8\pi^2 \sqrt{2N_c} f_V a_V^2 \exp \left[-a_V^2 \frac{\mathbf{k}_\perp^2}{\tau\bar{\tau}} \right]. \quad (6)$$

Here $\bar{\tau} = 1 - \tau$, f_V is the decay coupling constant and the a_V parameter determines the value of average transverse momentum of the quark in the vector meson. Generally, values of f_V, a_V are different for longitudinal and transverse polarization of the meson.

The subprocess amplitude is calculated within the MPA [5] where we keep the $k_\perp^2$ terms in the denominators of the amplitudes and the numerator of the TT amplitude. The gluonic corrections are treated in the form of the Sudakov factors which additionally suppress the end-point integration regions. The model leads to the following hierarchy of the amplitudes:

$$\text{LL: } \mathcal{M}_{0\nu,0\nu}^{V(g)} \propto 1; \quad \text{TL: } \mathcal{M}_{0\nu,+\nu}^{V(g)} \propto \frac{\sqrt{-t}}{Q}; \quad \text{TT: } \mathcal{M}_{+\nu,+\nu}^{V(g)} \propto \frac{k_\perp^2}{QM_V}. \quad (7)$$

Note, that the $L \rightarrow T$ and $T \rightarrow -T$ transitions are small and neglected in our analysis.

The GPD is a complicated function which depends on three variables. For the small momentum transfer $t \sim 0$ we use the double distribution form [10]

$$H^g(\bar{x}, \xi, t) = \left[\Theta(0 \leq \bar{x} \leq \xi) \int_{\frac{\bar{x}-\xi}{1-\xi}}^{\frac{\bar{x}+\xi}{1+\xi}} d\beta + \Theta(\xi \leq \bar{x} \leq 1) \int_{\frac{\bar{x}-\xi}{1-\xi}}^{\frac{\bar{x}+\xi}{1+\xi}} d\beta \right] \frac{\beta}{\xi} f(\beta, \alpha = \frac{\bar{x}-\beta}{\xi}, t) \quad (8)$$

with the simple factorizing ansatz for the double distributions $f(\beta, \alpha, t)$

$$f(\beta, \alpha, t \simeq 0) = g(\beta) \frac{3}{4} \frac{[(1-|\beta|)^2 - \alpha^2]}{(1-|\beta|)^3}. \quad (9)$$




Fig. 2. Model results for the GPD H^g in the small $\bar{x}$ range. The solid (dashed, dash-dotted) line represents the GPD at $\xi = 5$ (1 , 0.5) $\cdot 10^{-3}$

In this model, we calculate GPD [1] through the gluon distribution $g(\beta)$ and take the CTEQ5M results [11] for it. For the three ξ values the gluon GPD is shown in Fig. 2.

3 Cross section and spin observables

The t -dependence of the amplitudes is important in analyses of experimental data. For simplicity we parameterize it in the exponential form $M_{ii}(t) = M_{ii}(0) e^{t B_{ii}/2}$ for LL , TT , TL transitions. Experimentally, only the slope of the $\gamma^* p \rightarrow V p$ cross section is measured whereas information about individual slopes B_{ii} is absent. We suppose that $B_{LL} \sim B_{TT} \sim B_{TL}$. Estimations for the ρ meson production amplitudes are obtained using $f_{\rho L} = .216 \text{ GeV}$, $a_{\rho L} = 0.522 \text{ GeV}^{-1}$; $f_{\rho T} = .170 \text{ GeV}$; $a_{\rho T} = 0.65 \text{ GeV}^{-1}$.




Fig. 3a. The cross section for $\gamma^* p \rightarrow \rho^0 p$ vs. Q^2 for $W = 75 \text{ GeV}$.

Fig. 3b. The ratio of longitudinal and transverse cross sections for the ρ production vs. Q^2 at $W = 75 \text{ GeV}$ and $t = -0.15 \text{ GeV}^2$. Data are from H1 and ZEUS.

The cross section for the $\gamma^* p \rightarrow \rho p$ production integrated over t is shown in Fig. 3a. Good agreement with H1 and ZEUS experiments [6, 7] at HERA is to be observed. Note that the uncertainties in the gluon GPD provide about 20 – 30% errors in the cross section. The model results for the ratio of the cross section with longitudinal and transverse photon polarization R for the ρ production are shown in Fig. 3b. They are consistent with H1 and ZEUS experiments [6, 7].

In Fig. 4, we present our results for the three SDME at DESY energy range. Description of experimental data is reasonable. Our results for other SDME can be found in [1].

The A_{LL} asymmetry for a longitudinally polarized beam and target is sensitive to the polarized GPD. Really, the leading term in the A_{LL} asymmetry integrated over the azimuthal angle is determined through the interference between the unpolarized GPD H^g and the polarized $\tilde{H}^g$ distributions.

We expect a small value for the A_{LL} asymmetry at high energies because it is of the order of the ratio $\langle \tilde{H}^g \rangle / \langle H^g \rangle$ which is small at low x . At COMPASS energies $W = 10 \text{ GeV}$ the gluon contribution to the A_{LL} asymmetry of vector




Fig. 4 The Q^2 dependence of SDME on the ρ production at $t = -.15 \text{ GeV}^2$ and $W = 75 \text{ GeV}$. Data are taken from H1 and ZEUS.




Fig. 5. Gluon contribution to the A_{LL} asymmetry of vector meson production at $W = 5 \text{ GeV}$ (solid line) and $W = 10 \text{ GeV}$ (dashed line); $y \simeq 0.6$. The shaded bands reflect the uncertainties in the A_{LL} due to the error in $\tilde{H}^g \propto \Delta g$

meson production is small (Fig.5). At HERMES energies $W = 5 \text{ GeV}$ the major contribution to the amplitudes comes from the region $0.1 \leq \bar{x} \leq 0.2$ where $\Delta g/g$ is not small, which leads to a large value of the A_{LL} . The asymmetry is shown in Fig.5 together with the uncertainties in our predictions due to the error in the polarized gluon distribution. Note that our predictions should be valid for the ϕ production. In the case of ρ production the polarized quark GPD should be essential in the A_{LL} asymmetry in the low energy range.

4 Conclusion or Summary

Light vector meson electroproduction at small x was analyzed here within the GPD approach. The amplitudes were calculated using the MPA with the wave function dependent on the transverse quark momentum. The transverse quark momentum regularizes the end-point singularities in the amplitudes with transversally polarized photons. This gives a possibility to study spin effects in vector meson production. Within the GPD approach we find a fine description of the Q^2 dependence

of the cross section, R ratio and SDME for the ρ meson production at HERA energies. More results for ρ and ϕ production can be found in [1]. Some comparison with the two-gluon exchange model [12] can be found in [1] too.

We would like to point out that study of SDME gives important information on different $\gamma \rightarrow V$ hard amplitudes. By analyses of t dependencies of SDMEs the diffraction peak slopes of the TT and TL amplitudes can be defined. Unfortunately, the data on spin observables have large experimental errors. The new experimental results with reduced errors in SDME are extremely important. Information about polarized gluon distribution can be obtained from the A_{LL} asymmetry on ϕ meson production at HERMES and COMPASS energies which is sensitive to ΔG .

Thus, we can conclude that the vector meson photoproduction at small x is an excellent tool to probe the gluon GPD.

This work is supported in part by the Russian Foundation for Basic Research, Grant 06-02-16215 and by the Heisenberg-Landau program.

References

- [1] S.V. Goloskokov, P. Kroll: Euro. Phys. J. C**42** (2005) 281.
- [2] X. Ji: Phys. Rev. D**55** (1997) 7114;
A.V. Radyushkin: Phys. Lett. B**380** (1996) 417;
J.C. Collins: *et al.*, Phys. Rev. D**56** (1997) 2982.
- [3] L. Mankiewicz, G. Piller and T. Weigl: Eur. Phys. J. C**5** (1998) 119.
- [4] L. Mankiewicz and G. Piller: Phys. Rev. D**61** (2000) 074013;
I.V. Anikin and O.V. Teryaev: Phys. Lett. B**554** (2003) 51.
- [5] J. Botts and G. Sterman: Nucl. Phys. B**325** (1989) 62.
- [6] C. Adloff et al. [H1 Collaboration]: Eur. Phys. J. C**13** (2000) 371;
S. Aid et al. [H1 Collaboration]: Nucl. Phys. B**468** (1996) 3.
- [7] J. Breitweg et al. [ZEUS Collaboration]: Eur. Phys. J. C**6** (1999) 603;
S. Chekanov et al. [ZEUS Collaboration]: Nucl.Phys. B**718** (2005) 3.
- [8] S.V. Goloskokov, P. Kroll: Proc. of 16th International Spin Physics Symposium, Trieste, 2004, World Scientific, 2005, 421.
- [9] J. Bolz, J.G. Körner and P. Kroll: Z. Phys. A**350**(1994) 145.
- [10] I. V. Musatov and A. V. Radyushkin: Phys. Rev. D**61** (2000) 074027.
- [11] J. Pumplin, D. R. Stump, J. Huston, H. L. Lai, P. Nadolsky and W. K. Tung: JHEP **0207** (2002) 012.
- [12] S. J. Brodsky et al.: Phys. Rev. D **50** (1994) 3134.